

ISOSPECTRAL REDUCTION IN INFINITE GRAPHS

PEDRO DUARTE AND MARIA JOANA TORRES

ABSTRACT. L. A. Bunimovich and B. Z. Webb developed a theory for transforming a finite weighted graph while preserving its spectrum, referred as isospectral reduction theory. In this work we extend this theory to a class of operators on Banach spaces that include Markov type operators. We apply this theory to infinite countable weighted graphs admitting a finite structural set, a key concept in Bunimovich and Webb's theory.

Keywords: *Isospectral graph reduction, Markov operator, eigenvalue problem*

2000 Mathematics Subject Classification: 05C50, 47A75, 47D07

1. INTRODUCTION

L.A. Bunimovich and B.Z. Webb developed a theory for isospectral graph reduction in finite dimensional graphs (see [2, 1, 3]). This procedure maintains the spectrum of the graph's adjacency matrix up to a set of eigenvalues known beforehand from its graph structure. More precisely, the authors introduce a concept of transformation of a graph (either by reduction or expansion) that can be used to simplify the structure of a graph while preserving the eigenvalues of the graph's adjacency matrix. In order to not contradict the fundamental theorem of algebra, isospectral graph transformations preserve the spectrum of the graph (in particular the number of eigenvalues) by permitting edges to be weighted by functions of a spectral parameter λ (see [2, Theorem 3.5.]). Thus such transformations allow one to modify the topology of a network (changing the interactions, reducing or increasing the number of nodes), while maintaining properties related to the network's dynamics.

The work [2] contains also an application of this isospectral theory to a class of dynamical systems modeling dynamical networks. In particular, the authors provide a sufficient condition for a dynamical network having a globally attracting fixed point. In [1], the theory of isospectral reduction is also used to improve classical eigenvalues estimates.

More recently, in [4], we have proven that isospectral graph reductions also preserve the eigenvectors associated to the eigenvalues of the graph's weighted adjacency matrix. We explain how the isospectral reduction procedure can be used to efficiently update the eigenvector of a large sparse matrix when only a small number of its entries is modified.

As an application we propose an updating algorithm for the maximal eigenvector of the Markov matrix associated to a large sparse dynamical network.

Because our spectral approach to isospectral graph reduction theory is based on *eigenvectors*, instead of *eigenvalues*, it was a natural question to ask about possible generalizations of this theory to infinite dimensions.

We believe there are many possible such extensions to infinite dimensional models. In this work we develop a couple of abstract settings where such generalizations hold. In a future work we expect to develop isospectral reduction in a different setting more suitable for applications to discrete deterministic dynamical systems.

The theory applies to a class of bounded linear operators acting on spaces of L^1 -integrable functions. The operators considered are written as a sum of a diagonal plus a Markov operator. A key concept in Bunimovich-Webb's isospectral theory is that of a *structural set*. In this work we give three different concepts of structural sets (see Definitions 3.4, 3.5 and 3.6) and for each of them prove a corresponding isospectral theorem (see Theorems 3.12, 3.8 and 3.18).

An application of Theorem 3.18 is given to weighted countably infinite graphs with a finite structural set (see Theorem 4.2). We also propose a numerical algorithm to approximate the eigenfunctions of such weighted graphs. We conclude the manuscript with a concrete application of this method.

The paper is organized as follows:

In Section 2 we describe the isospectral graph reduction theory and the reduction statements for finite graphs.

In Section 3 we generalize the isospectral graph reduction theory to infinite dimensional models.

In Section 4 we apply the infinite dimension isospectral reduction theory, developed in Section 3, to countably infinite graphs with a finite structural set. We also propose a numerical algorithm to approximate the eigenfunctions of such graphs.

In Section 5 we present a concrete application of the algorithm proposed in Section 4.

2. ISOSPECTRAL REDUCTION IN FINITE GRAPHS

In this section we describe the isospectral graph reduction theory and the reduction statements for finite graphs.

Definition 2.1 (Finite weighted graph). *A finite weighted graph is a pair $G = (V, w)$ where V is a finite set and $w : V \times V \rightarrow \mathbb{C}$ is any function, called the weight function of G . We denote by $\mathcal{A} = \mathcal{A}_w : \mathbb{C}^V \rightarrow \mathbb{C}^V$ the operator defined by the weighted adjacency matrix $(w(v, v'))_{v, v' \in V}$.*

A *path* $\gamma = (v_0, \dots, v_p)$ in the graph $G = (V, w)$ is an ordered sequence of vertices $v_0, \dots, v_p \in V$ such that $w(v_\ell, v_{\ell+1}) \neq 0$ for $0 \leq \ell \leq p-1$. The integer p is called the

length of γ . If the vertices $v_0, \dots, v_{p-1}$ are all distinct the path γ is called *simple*. If $v_0 = v_p$ then γ is called a *closed path*. A closed path of length 1 is called a *loop*. Finally, we call *cycle* any simple closed path.

If $S \subseteq V$ we will write $S^c = V \setminus S$.

Definition 2.2 (Structural set). *Let $G = (V, w)$. A nonempty vertex set $S \subseteq V$ is a structural set for G if each cycle of G , that is not a loop, contains a vertex in S .*

Given a structural set S , we call *branch of (G, S)* any simple path $\beta = (v_0, v_1, \dots, v_{p-1}, v_p)$ such that $v_1, \dots, v_{p-1} \in S^c$ and $v_0, v_p \in V$. We denote by $\mathcal{B} = \mathcal{B}_{G,S}$ the set of all branches of (G, S) . Given vertices $v, v' \in V$, we denote by $\mathcal{B}_{vv'}$ the set of all branches in $\mathcal{B}$ that start in v and end in v' . Define $\Sigma = \{w(v, v) : v \in S^c\}$ and let $\lambda \in \mathbb{C} \setminus \Sigma$. For each branch $\beta = (v_0, v_1, \dots, v_p)$ we define the λ -weight of β as follows:

$$w(\beta, \lambda) := w(v_0, v_1) \prod_{\ell=1}^{p-1} \frac{w(v_\ell, v_{\ell+1})}{\lambda - w(v_\ell, v_\ell)}. \quad (2.1)$$

Given $v, v' \in V$ set

$$R_{S,\lambda}(v, v') := \sum_{\beta \in \mathcal{B}_{vv'}} w(\beta, \lambda). \quad (2.2)$$

The *reduced operator* $\mathcal{R}_S(\lambda) : \mathbb{C}^S \rightarrow \mathbb{C}^S$ is the operator with matrix $(R_{S,\lambda}(v, v'))_{v,v' \in S}$.

In [2] the reduced operator $\mathcal{R}_S$ is viewed as a matrix indexed in $S \times S$ with values in the field $\mathbb{W}[\lambda]$ of all rational functions $f(\lambda) = \frac{p(\lambda)}{q(\lambda)}$, where $p(\lambda)$ and $q(\lambda)$ are polynomials. In their treatment Bunimovich and Webb consider, more generally, weighted adjacency matrices $\mathcal{A}$ with values in the field $\mathbb{W}[\lambda]$ instead of $\mathbb{C}$, so that the reduced matrix $\mathcal{R}_S$ lives in the same space of $\mathbb{W}[\lambda]$ -valued matrices. Given a matrix $\mathcal{A}(\lambda) \in \mathbb{W}[\lambda]^{V \times V}$ its spectrum is defined in [2, Definition 3.1] by $\text{sp}(\mathcal{A}(\lambda)) = P \setminus Q$ where $P = \{\lambda \in \mathbb{C} : p(\lambda) = 0\}$, $Q = \{\lambda \in \mathbb{C} : q(\lambda) = 0\}$ and $\det(\mathcal{A}(\lambda) - \lambda I) = p(\lambda)/q(\lambda)$. In the context of the previous definitions, starting with a complex valued matrix $\mathcal{A} \in \mathbb{C}^{V \times V}$, by [2, Corollary 3] the spectrum of the reduced operator $\mathcal{R}_S(\lambda)$ matches the following definition, which is more suitable for our infinite dimensional isospectral reduction.

Definition 2.3. *We define the spectrum of the family of operators $\mathcal{R}_S(\lambda)$, denoted by $\text{sp}(\mathcal{R}_S)$, to be*

$$\text{sp}(\mathcal{R}_S) := \{\lambda \in \mathbb{C} \setminus \Sigma : \det(\mathcal{R}_S(\lambda) - \lambda I) = 0\}.$$

A simplified¹ version of Bunimovich-Webb isospectral reduction theorem (see [2, Theorem 3.5.]) can be stated as follows:

¹This statement corresponds to [2, Corollary 3] where the adjacency matrix has complex entries.

Theorem 1 (Bunimovich-Webb). *Given a structural set S for a graph $G = (V, w)$,*

$$\text{sp}(\mathcal{A}) \setminus \Sigma = \text{sp}(\mathcal{R}_S).$$

We have stated our reduction results in [4, Theorem 1, Proposition 2.1] in terms of restriction and extensions of eigenvectors. The following theorem states that isospectral graph reduction preserves the eigenvectors associated to eigenvalues of the graph's weighted adjacency matrix.

Theorem 2 ([4, Theorem 1]). *Given a graph $G = (V, w)$, let $\lambda_0 \in \mathbb{C} \setminus \Sigma$ be an eigenvalue of $\mathcal{A} : \mathbb{C}^V \rightarrow \mathbb{C}^V$ and $u = (u_1, u_2, \dots, u_n) \in \mathbb{C}^n$ be the corresponding eigenvector, $\mathcal{A}u = \lambda_0 u$. Assume that $S = \{m+1, \dots, n\}$ is a structural set for G . Then λ_0 is also an eigenvalue of $\mathcal{R}_S(\lambda_0)$ and $\mathcal{R}_S(\lambda_0)u_S = \lambda_0 u_S$, where $u_S = (u_{m+1}, \dots, u_n)$ is the restriction of u to S .*

To explain how to reconstruct the eigenvector of $\mathcal{A}$ from the eigenvector of the reduced matrix $\mathcal{R}_S(\lambda_0)$ we need the following concept of depth of vertex $v \in V$.

Definition 2.4 (Depth of a vertex). *The depth of a vertex $v \in V$ is defined recursively as follows.*

- (1) *A vertex $v \in S$ has depth 0.*
- (2) *A vertex $v \in S^c$ has depth n iff v has no depth less than n , and $w(v, v') \neq 0$ implies v' has depth $< n$, for all $v' \in V$.*

We denote by S_n the set of all vertices of depth $\leq n$. Because S is a structural set, every vertex v has a finite depth.

If $\lambda_0 \in \mathbb{C} \setminus \Sigma$ is an eigenvalue of $\mathcal{A}$, by Theorem 2 it is also an eigenvalue of the reduced matrix $\mathcal{R}_S(\lambda_0)$. Knowing the eigenvector u_S of this reduced matrix, we can recover the corresponding eigenvector of $\mathcal{A}$ as follows:

Proposition 2.1 ([4, Proposition 2.1]). *If $\lambda_0 \in \mathbb{C} \setminus \Sigma$ is an eigenvalue of $\mathcal{A}$ and $u_S = (u_v^S)_{v \in S}$ is an eigenvector of the reduced matrix $\mathcal{R}_S(\lambda_0)$ then the following recursive relations*

$$\left\{ \begin{array}{ll} u_v = u_v^S & \text{for } v \in S_0 = S \\ u_\ell = \sum_{v' \in S_{n-1}} \frac{w(\ell, v')}{\lambda_0 - w(\ell, \ell)} u_{v'} & \text{for all } \ell \in S_n \setminus S_{n-1} \end{array} \right. \quad (2.3)$$

uniquely determine an eigenvector u of $\mathcal{A}$ associated to λ_0 .

Denote by $\Pi_S : \mathbb{C}^V \rightarrow \mathbb{C}^S$ the S -restriction projection, and let $\Phi_S : \mathbb{C} \setminus \Sigma \rightarrow \text{Mat}_{V \times S}(\mathbb{C})$ be the function that to each $\lambda \in \mathbb{C} \setminus \Sigma$ associates the *reconstruction operator* $\Phi_S(\lambda) : \mathbb{C}^S \rightarrow \mathbb{C}^V$ where $u = \Phi_S(\lambda)w$ is recursively defined by

$$\begin{cases} u_v = w_v & \text{for } v \in S_0 = S \\ u_\ell = \sum_{v' \in S_{n-1}} \frac{w(\ell, v')}{\lambda_0 - w(\ell, \ell)} u_{v'} & \text{for all } \ell \in S_n \setminus S_{n-1} \end{cases}. \quad (2.4)$$

These maps are inverse of each other in the sense that $\Pi_S \circ \Phi_S(\lambda) = \text{id}_{\mathbb{C}^S}$ for all $\lambda \in \mathbb{C} \setminus \Sigma$. Finally notice that the reconstruction operator $\Phi_S(\lambda)$ is analytic in $\lambda \in \mathbb{C} \setminus \Sigma$.

The aim of this paper is to extend the reduction statements in Theorem 1, Theorem 2 and Proposition 2.1 to infinite dimensional models.

3. ISOSPECTRAL REDUCTION IN INFINITE DIMENSIONAL MODELS

In this section we generalize the isospectral graph reduction theory to infinite dimensional models.

Our infinite dimensional model (V, K, d, S) is defined by a measurable space $(V, \mathcal{F})$, a kernel K on V , a bounded measurable function $d : V \rightarrow \mathbb{C}$, and a structural set $S \subseteq V$.

Definition 3.1 (Kernel). *A kernel on V is any function $K : V \times \mathcal{F} \rightarrow \mathbb{C}$ such that*

- (1) *the function $B \mapsto K(x, B)$, from $\mathcal{F}$ to $\mathbb{C}$, is a complex measure for any $x \in V$;*
- (2) *the function $x \mapsto K(x, B)$, from V to $\mathbb{C}$, is $\mathcal{F}$ -measurable for any $B \in \mathcal{F}$.*

In particular a kernel K determines a function $K : V \rightarrow \mathcal{M}(V, \mathbb{C})$ that to each $x \in V$ associates the measure K_x . The notation $\mathcal{M}(V, \mathbb{C})$ stands for the space of complex measures on $(V, \mathcal{F})$. We use the following notation for the integral of an $\mathcal{F}$ -measurable function $f : V \rightarrow \mathbb{C}$ w.r.t. K_x

$$\int_V f dK_x = \int_V f(y) K(x, dy).$$

We also define the positive kernel $|K| : V \times \mathcal{F} \rightarrow [0, +\infty]$

$$|K|(x, B) = |K_x|(B),$$

where $|K_x|(B)$ stands for the total variation of K_x on B .

Definition 3.2. *We say that a kernel K has no diagonal part on a set $B \in \mathcal{F}$ when for all $z \in B$, $K(z, \{z\}) = 0$.*

Let us fix a positive finite reference measure μ on $(V, \mathcal{F})$.

Definition 3.3 (μ -bounded kernel). *We say that a kernel K is μ -bounded when there exists $M < +\infty$ such that for all $x \in V$ and $B \in \mathcal{F}$, $|K(x, B)| \leq M \mu(B)$.*

This means that K_x is absolutely continuous with respect to μ , i.e., $K(x, dy) = h(x, y) \mu(dy)$ for some measurable function $h : V \times V \rightarrow \mathbb{R}$.

Given a Banach space $(\mathcal{B}, \|\cdot\|)$, we denote by $\mathcal{L}(\mathcal{B})$ the Banach algebra of bounded linear operators on $\mathcal{B}$. Given $Q \in \mathcal{L}(\mathcal{B})$, the *operator norm* of Q is defined by $\|Q\| := \sup_{\|f\|=1} \|Qf\|$. Let $(L^1(V), \|\cdot\|_{L^1(V)})$ denote the Banach space of complex μ -integrable functions with the usual L^1 norm $\|f\|_{L^1(V)} := \int_V |f| d\mu$, and let $(L^\infty(V), \|\cdot\|_\infty)$ denote the Banach algebra of complex bounded $\mathcal{F}$ -measurable functions with the usual sup norm $\|f\|_\infty := \sup_{x \in V} |f(x)|$.

Given a μ -bounded kernel K and a measurable function $d \in L^\infty(V)$, consider the operator $\mathcal{A} = \mathcal{A}_{d,K} : L^1(V) \rightarrow L^1(V)$,

$$(\mathcal{A}f)(x) := d(x)f(x) + \int_V f(y) K(x, dy). \quad (3.1)$$

When $d \equiv 0$, kernel K determines an operator $\mathcal{Q} = \mathcal{Q}_K : L^1(V) \rightarrow L^1(V)$,

$$(\mathcal{Q}f)(x) := \int_V f(y) K(x, dy), \quad (3.2)$$

that we will refer as the *Markov operator* with kernel K .

The following proposition is a simple observation.

Proposition 3.1. *If K is μ -bounded and $d \in L^\infty(V)$ then $\mathcal{A} \in \mathcal{L}(L^1(V))$. In particular also $\mathcal{Q} \in \mathcal{L}(L^1(V))$.*

We make the following assumptions throughout the rest of this section.

- (A1) K is a μ -bounded kernel on V w.r.t. a positive finite measure μ on $(V, \mathcal{F})$.
- (A2) $d : V \rightarrow \mathbb{C}$ is a bounded measurable function.

Let $S \subseteq V$ (in the sequel S will be a structural set, in a sense to be defined below) and let $\Sigma_d := \overline{d(V \setminus S)}$ (where the overline means topological closure in $\mathbb{C}$). Consider the σ -algebra $\mathcal{F}_S := \{S \cap B : B \in \mathcal{F}\}$, the restricted measure $\mu_S : \mathcal{F}_S \rightarrow [0, +\infty[$ defined by $\mu_S(B) = \mu(B)$, and let $L^1(S)$ denote the space of complex μ_S -integrable functions w.r.t. μ_S .

Given $\lambda \in \mathbb{C} \setminus \Sigma_d$, we define the *reduced operator* $\mathcal{R}_S(\lambda) = \mathcal{R}_{S,d,K}(\lambda) : L^1(S) \rightarrow L^1(S)$ by

$$(\mathcal{R}_S(\lambda) f)(x) := d(x)f(x) + \int_S f(y) R_{S,\lambda}(x, dy) , \quad (3.3)$$

where, for $B \in \mathcal{F}_S$,

$$R_{S,\lambda}(x, B) := \sum_{n=1}^{\infty} K_{S,\lambda}^{(n)}(x, B) , \quad (3.4)$$

with $K_{S,\lambda}^{(1)}(x, B) := K(x, B)$ and for $n \geq 2$,

$$K_{S,\lambda}^{(n)}(x, B) := \int_{S^c} \cdots \int_{S^c} \frac{K(x, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, B)}{\prod_{p=1}^{n-1} (\lambda - d(z_p))} .$$

This is a formal definition. The reduced operator may not be well defined if the series (3.4) fails to converge. We will now define two concepts of structural set for which the reduced operator becomes well defined.

Let $S \subseteq V$. Given $x \in V$, $B \in \mathcal{F}$ and $n \geq 2$ define the *S-taboo measure*

$$\tau_{S,n}(x, B) = \tau_{S,n,K}(x, B) := \int_{S^c} \cdots \int_{S^c} K(x, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, B).$$

Let us say that a sequence $t_n > 0$ converges to 0 *super exponentially* when

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log t_n = -\infty.$$

Definition 3.4 (Structural set of type A). *A nonempty set $S \subseteq V$ is called a structural set of type A for K if and only if there exists a sequence t_n converging to 0 super exponentially such that $|\tau_{S,n}(x, B)| \leq t_n \mu(B)$ for all $x \in V$, $B \in \mathcal{F}$ and $n \geq 2$.*

We call point-set map on V to any map $F : V \rightarrow \mathcal{P}(V)$, where $\mathcal{P}(V)$ stands for the power set of V . We write $F : V \rightrightarrows V$ to express that F is a point-set map on V . We define recursively for all $n \geq 1$ and $x \in V$,

$$\varphi^n(x) = \cup \{ \varphi(y) : y \in \varphi^{n-1}(x) \}.$$

Definition 3.5 (Structural set of type B). *A nonempty set $S \subseteq V$ is called a structural set of type B for K if and only if defining the point-set map $\varphi_{S^c} : S^c \rightrightarrows S^c$,*

$$\varphi_{S^c}(x) := S^c \cap \text{supp}(K_x)$$

one has

- (1) for all $x \in S^c$, there exists $n \in \mathbb{N}$ such that $(\varphi_{S^c})^n(x) = \emptyset$,

(2) setting $n_S : S^c \rightarrow \mathbb{N}$, $n_S(x) := \min\{k \in \mathbb{N} : (\varphi_{S^c})^k(x) = \emptyset\}$ then,

$$\int_{S^c} n_S(x) \mu(dx) < +\infty.$$

Remark 3.1. *The concepts of structural sets of type A and B are logically independent.*

Remark 3.2. *If (V, K) admits a structural set of type A or B then K has no diagonal part on S^c . Indeed, for structural sets of type A, notice that if $p := |K(z, \{z\})| > 0$ then $|\tau_{S,n}(z, \{z\})| \geq p^n$ for all $n \in \mathbb{N}$. For structural sets of type B, it follows from Definition 3.5 (1) that K has no diagonal part on S^c .*

Remark 3.3. *If V is finite and S is a structural set of type A or B for K then S is a structural set in the sense of Definition 2.2.*

From (A1) it follows that each measure K_x has a density $h = dK_x/d\mu \in L^1(V)$ with $\|dK_x/d\mu\|_\infty \leq M$. We define

$$\|K\|_\infty := \sup_{x \in V} \left\| \frac{dK_x}{d\mu} \right\|_\infty. \quad (3.5)$$

We use this norm to measure distances between kernels.

Definition 3.6 (Structural set of type A quasi-B). *A nonempty set $S \subseteq V$ is called a structural set of type A quasi-B for K if and only if there exists a sequence of μ -bounded kernels K_n , $n \in \mathbb{N}$, such that*

- (1) *S is a structural set of type A for K ;*
- (2) $\lim_{n \rightarrow +\infty} \|K - K_n\|_\infty = 0$;
- (3) *S is a structural set of type B for K_n , for all $n \in \mathbb{N}$.*

Proposition 3.2. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type A quasi-B for K . Then $\lim_{n \rightarrow +\infty} \mathcal{A}_{d, K_n} = \mathcal{A}_{d, K}$ in $\mathcal{L}(L^1(V))$.*

Proof. Let $K(x, dy) = h(x, y) \mu(dy)$ and $K_n(x, dy) = h_n(x, y) \mu(dy)$. Given $f \in L^1(V)$ we have that

$$\begin{aligned}
\|\mathcal{A}_{d,K}f - \mathcal{A}_{d,K_n}f\|_{L^1(V)} &= \int_V |(\mathcal{A}_{d,K}f - \mathcal{A}_{d,K_n}f)(x)| \mu(dx) \\
&= \int_V \left| \int_V (K(x, dy) - K_n(x, dy))f(y) \right| \mu(dx) \\
&= \int_V \left| \int_V (h(x, y) - h_n(x, y))f(y) \mu(dy) \right| \mu(dx) \\
&\leq \int_V \int_V |h(x, y) - h_n(x, y)| |f(y)| \mu(dy) \mu(dx) \\
&\leq \mu(V) \|h - h_n\|_\infty \|f\|_{L^1(V)}.
\end{aligned}$$

By Definition 3.6 (2) (of type A quasi-B structural sets) we have that $\|h - h_n\|_\infty \rightarrow 0$. Thus, $\lim_{n \rightarrow +\infty} \mathcal{A}_{d,K_n} = \mathcal{A}_{d,K}$ in $\mathcal{L}(L^1(V))$. $\square$

3.1. Reduced Operator. We shall now prove that for structural sets of type A or type B, the reduced operators $\mathcal{R}_S(\lambda)$ defined by (3.3) are well defined, bounded and, moreover, that the function $\mathcal{R}_S : \mathbb{C} \setminus \Sigma_d \rightarrow \mathcal{L}(L^1(S))$ is analytic.

Lemma 3.3. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type A. Given $\lambda \in \mathbb{C} \setminus \Sigma_d$, the kernel series $\sum_{n=1}^\infty K_{S,\lambda}^{(n)}(x, B)$ converges absolutely and uniformly on $S \times \mathcal{F}_S$. Moreover, the operators $\mathcal{R}_S(\lambda)$ defined by (3.3) are bounded.*

Proof. For each $r > 0$, define the open set $\Omega_r := \{\lambda \in \mathbb{C} : \text{dist}(\lambda, \Sigma_d) > r\}$, so that $\mathbb{C} \setminus \Sigma_d = \cup_{r>0} \Omega_r$. Given $\lambda \in \Omega_r$ and a list of points $\bar{z} = (z_1, \dots, z_{n-1}) \in V^{n-1}$, the analytic function

$$f_{\bar{z}}(\lambda) := \frac{1}{\prod_{p=1}^{n-1} (\lambda - d(z_p))}$$

is bounded by $r^{-(n-1)}$. Therefore, given $x \in S$ and $B \in \mathcal{F}_S$

$$|K_{S,\lambda}^{(n)}(x, B)| \leq \frac{|\tau_{S,n}(x, B)|}{r^{n-1}} \leq \frac{t_n}{r^{n-1}} \mu(B).$$

But since the sequence $t_n \searrow 0$ super exponentially, applying d'Alembert's criterion (ratio test) we can conclude that

$$C_r := \sum_{n=1}^\infty \frac{t_n}{r^{n-1}} < +\infty.$$

From the previous bound on $|K_{S,\lambda}^{(n)}(x, B)|$ we infer that for all $f \in L^1(S)$

$$\int \left| \int f(y) K_{S,\lambda}^{(n)}(x, dy) \right| d\mu(x) \leq \frac{t_n}{r^{n-1}} \|f\|_{L^1(S)} \mu(S).$$

Hence the Markov operator defined by $K_{S,\lambda}^{(n)}$ is in $\mathcal{L}(L^1(S))$ with norm bounded by $\frac{t_n}{r^{n-1}} \mu(S)$. A straightforward calculation shows that the reduced operator $\mathcal{R}_S(\lambda)$ has norm $\|\mathcal{R}_S(\lambda)\|_{L^1(S)} \leq \|d\|_\infty + C_r \mu(S)$, for $\lambda \in \Omega_r$. $\square$

Proposition 3.4. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type A. Then the function $\mathcal{R}_S : \mathbb{C} \setminus \Sigma_d \rightarrow \mathcal{L}(L^1(S))$ is analytic.*

Proof. Consider the open sets Ω_r , $r > 0$, and the analytic function $f_{\bar{z}}(\lambda)$, $\bar{z} \in V^{n-1}$, introduced in the proof of Lemma 3.3. Differentiating $f_{\bar{z}}(\lambda)$ in λ we get

$$f'_{\bar{z}}(\lambda) = - \left(\prod_{p=1}^{n-1} \frac{1}{\lambda - d(z_p)} \right) \sum_{p=1}^{n-1} \frac{1}{\lambda - d(z_p)},$$

with $|f'_{\bar{z}}(\lambda)| \leq n r^{-n}$, for all $\lambda \in \Omega_r$.

For each $n \in \mathbb{N}$ consider the operator $\mathcal{K}_n(\lambda) : L^1(S) \rightarrow L^1(S)$ defined by

$$\begin{aligned} (\mathcal{K}_n(\lambda) h)(x) &:= \int_S h(y) K_{S,\lambda}^{(n)}(x, dy) \\ &= \int_S \int_{S^c} \cdots \int_{S^c} h(y) f_{z_1, \dots, z_{n-1}}(\lambda) K(x, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, dy). \end{aligned}$$

By the previous bound, arguing as in Lemma 3.3, these operators are analytic with

$$\left\| \frac{d}{d\lambda} \mathcal{K}_n(\lambda) \right\| \leq n \frac{t_n}{r^n} \mu(S),$$

for all $\lambda \in \Omega_r$. Thus, by d'Alembert's criterion, the series of analytic functions $\lambda \mapsto \sum_{n=1}^{\infty} \frac{d}{d\lambda} \mathcal{K}_n(\lambda) \in \mathcal{L}(L^1(S))$ converges uniformly on Ω_r . This proves that $\lambda \mapsto \mathcal{R}_S(\lambda)$ is analytic on $\mathbb{C} \setminus \Sigma_d$. $\square$

Lemma 3.5. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type B. Given $\lambda \in \mathbb{C} \setminus \Sigma_d$, the kernel series $\sum_{n=1}^{\infty} K_{S,\lambda}^{(n)}(x, B)$ converges absolutely and uniformly on $S \times \mathcal{F}_S$. Moreover, the operators $\mathcal{R}_S(\lambda)$ defined by (3.3) are bounded.*

Proof. Define $D_n = \{z \in S^c : n_S(z) \geq n\}$, where $n_S(z)$ is the function introduced in Definition 3.5(2). Given $B \in \mathcal{F}_S$ and $x \in S$ one has

$$K_{S,\lambda}^{(n)}(x, B) = \int_{D_1} \cdots \int_{D_{n-1}} \frac{K(x, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, B)}{\prod_{p=1}^{n-1} (\lambda - d(z_p))}.$$

Consider the open sets Ω_r , $r > 0$, introduced in the proof of Lemma 3.3. From (A1), arguing as in the proof of Lemma 3.3, for all $\lambda \in \Omega_r$ we have that

$$|K_{S,\lambda}^{(n)}(x, B)| \leq \frac{M^n}{r^{n-1}} \mu(D_1) \cdots \mu(D_{n-1}) \mu(B). \quad (3.6)$$

But since

$$\sum_{n=1}^{\infty} \mu(D_n) = \int_{S^c} n_S(x) \mu(dx) < +\infty,$$

applying d'Alembert's criterion (ratio test) we conclude that

$$C_r := \sum_{n=1}^{\infty} \frac{M^n}{r^{n-1}} \mu(D_1) \cdots \mu(D_{n-1}) < +\infty.$$

A straightforward calculation shows that the operator $\mathcal{R}_S(\lambda) : L^1(S) \rightarrow L^1(S)$ has norm $\|\mathcal{R}_S(\lambda)\|_{L^1(S)} \leq \|d\|_{\infty} + C_r \mu(S)$, for $\lambda \in \Omega_r$. $\square$

Proposition 3.6. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type B. Then the function $\mathcal{R}_S : \mathbb{C} \setminus \Sigma_d \rightarrow \mathcal{L}(L^1(S))$ is analytic.*

Proof. This proof is very similar to the one of Proposition 3.4. Consider the open sets Ω_r , $r > 0$, introduced in the proof of Lemma 3.3 and the operators $\mathcal{K}_n(\lambda)$, $n \in \mathbb{N}$, introduced in the proof of Proposition 3.4. By the previous bound in (3.6), arguing as in the proof of Proposition 3.4, these operators are analytic with

$$\left\| \frac{d}{d\lambda} \mathcal{K}_n(\lambda) \right\| \leq n \frac{M^n}{r^n} \mu(D_1) \cdots \mu(D_{n-1}) \mu(S),$$

for all $\lambda \in \Omega_r$. Thus, by d'Alembert's criterion, the series of analytic functions $\lambda \mapsto \sum_{n=1}^{\infty} \frac{d}{d\lambda} \mathcal{K}_n(\lambda) \in \mathcal{L}(L^1(S))$ converges uniformly on Ω_r . This proves that $\lambda \mapsto \mathcal{R}_S(\lambda)$ is analytic on $\mathbb{C} \setminus \Sigma_d$. $\square$

3.2. Type B isospectral reduction. In this subsection we prove a isospectral reduction theorem for structural sets of type B.

For such structural sets the following key definition captures the idea of depth of a point in V (see Definition 2.4).

Definition 3.7. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type B. We define, recursively in $n \in \mathbb{N}$, a measurable subset $S_n \subseteq V$ which can be regarded as the set of states $x \in V$ of depth $\leq n$.*

- (1) $S_0 := S$,
- (2) $S_n := \{x \in V : |K|(x, V \setminus S_{n-1}) = 0\} \cup S_{n-1}$.

Proposition 3.7. *The sets S_n cover V , i.e., $V = \cup_{n \geq 0} S_n$.*

Proof. Given $x \in V$, let $n = n_S(x)$ be as in Definition 3.5 (2). We will prove by induction that $\varphi_{S^c}^{n-i}(x) \subseteq S_i$ for all $i = 1, \dots, n$. Then, taking $i = n$, this shows that $\{x\} = \varphi_{S^c}^0(x) \subseteq S_n$.

Let $y \in \varphi_{S^c}^{n-1}(x)$. Then $\varphi_{S^c}(y) = \text{supp}(K_y) \cap S^c = \emptyset$, which implies $|K|(y, S^c) = 0$, and hence $|K|(y, V \setminus S_0) = |K|(y, S^c) = 0$. Thus $y \in S_1$, which proves the claim for $i = 1$.

Assume, by induction hypothesis, that $\varphi_{S^c}^{n-i+1}(x) \subseteq S_{i-1}$, and let $y \in \varphi_{S^c}^{n-i}(x)$, which implies that $\varphi_{S^c}(y) \subseteq \varphi_{S^c}^{n-i+1}(x) \subseteq S_{i-1}$.

We consider two cases:

First assume that $\mu(\varphi_{S^c}(y)) = 0$. In this case, because K_y is absolutely continuous w.r.t. μ , we have that $|K|(y, S^c) = 0$. Thus $|K|(y, V \setminus S_0) = |K|(y, S^c) = 0$, and consequently $y \in S_1 \subseteq S_i$.

Assume now that $\mu(\varphi_{S^c}(y)) > 0$. Since $|K(y, z)| > 0$ for μ -a.e. $z \in \text{supp}(K_y)$, and $\varphi_{S^c}(y) \subseteq S_{i-1}$, we have that $\text{supp}(K_y) \subseteq S_{i-1}$ and so $|K|(y, V \setminus S_{i-1}) = 0$. Therefore $y \in S_i$.

This proves that $\varphi_{S^c}^{n-i}(x) \subseteq S_i$, and concludes the inductive argument. $\square$

Given $u \in L^1(V)$, we define the functions u_S and u_{S^c} in $L^1(V)$,

$$u_S(x) := \begin{cases} u(x) & \text{if } x \in S \\ 0 & \text{if } x \in S^c \end{cases} \quad u_{S^c}(x) := \begin{cases} u(x) & \text{if } x \in S^c \\ 0 & \text{if } x \in S \end{cases}.$$

From now on, whenever appropriate we will identify u_S with a function in $L^1(S)$.

Definition 3.8. *The spectrum of the family of operators $\mathcal{R}_S(\lambda)$, denoted by $\text{sp}(\mathcal{R}_S)$, is the set of $\lambda \in \mathbb{C} \setminus \Sigma_d$ such that $\mathcal{R}_S(\lambda) - \lambda I$ is a non invertible operator on $L^1(S)$.*

Theorem 3.8. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type B. Then*

- (1) $\text{sp}(\mathcal{A}) \setminus \Sigma_d \subseteq \text{sp}(\mathcal{R}_S)$.
- (2) Given $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$, λ_0 is an eigenvalue of $\mathcal{A}$ iff λ_0 is an eigenvalue of $\mathcal{R}_S(\lambda_0)$.
- (3) If $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ is an eigenvalue of $\mathcal{A}$ and $u \in L^1(V)$ is an associated eigenfunction, $\mathcal{A}u = \lambda_0 u$, then $\mathcal{R}_S(\lambda_0)u_S = \lambda_0 u_S$, i.e., u_S is the corresponding eigenfunction for $\mathcal{R}_S(\lambda_0)$.
- (4) If $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ is an eigenvalue of $\mathcal{R}_S(\lambda_0)$ and v is an associated eigenfunction, $\mathcal{R}_S(\lambda_0)v = \lambda_0 v$, then the following recursive relations

$$\begin{cases} u(x) = v(x) & \text{for } x \in S_0 = S \\ u(z) = \int_{S_{n-1}} \frac{K(z, dy)}{\lambda_0 - d(z)} u(y) & \text{for all } z \in S_n \setminus S_{n-1} \end{cases} \quad (3.7)$$

uniquely determine an eigenfunction $u \in L^1(V)$ such that $\mathcal{A}u = \lambda_0 u$.

Remark 3.4. Under the assumptions of Theorem 3.8, if S is finite then the equality holds in item (1), i.e., $\text{sp}(\mathcal{A}) \setminus \Sigma_d = \text{sp}(\mathcal{R}_S)$. Indeed, in this case, the reduced operator is finite dimensional and, consequently, the equality follows from item (2).

To prove Theorem 3.8 we need the following lemma.

Lemma 3.9. Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type B. Given $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$, $f \in L^1(V)$ and $v \in L^1(S)$, defining recursively a function $u : V \rightarrow \mathbb{C}$ by

$$\begin{cases} u(x) = v(x) & \text{for } x \in S_0 = S \\ u(z) = -\frac{f(z)}{\lambda_0 - d(z)} + \int_{S_{n-1}} \frac{K(z, dy)}{\lambda_0 - d(z)} u(y) & \text{for all } z \in S_n \setminus S_{n-1} \end{cases} \quad (3.8)$$

then $u \in L^1(V)$. Furthermore, $\|u\|_{L^1(V)} \leq C(\|f\|_{L^1(V)} + \|v\|_{L^1(S)})$, for some constant $C = C(\lambda_0) > 0$.

Given $B \in \mathcal{F}$, let us denote by 1_B the indicator function of B .

Given a function $v : S \rightarrow \mathbb{C}$ we denote by $\bar{v}$ the extension $\bar{v} : V \rightarrow \mathbb{C}$,

$$\bar{v}(x) = \begin{cases} v(x) & \text{if } x \in S \\ 0 & \text{if } x \notin S \end{cases}.$$

Given $\lambda \in \mathbb{C} \setminus \Sigma_d$, we introduce the operator $\mathcal{D}(\lambda) : L^1(V) \rightarrow L^1(V)$ defined by

$$(\mathcal{D}(\lambda)h)(z) := 1_{S^c}(z) \frac{h(z)}{\lambda - d(z)}, \quad (3.9)$$

which clearly is a bounded operator.

To simplify notations we will write $\mathcal{Q}$ instead of $\mathcal{Q}_K$, and $\mathcal{D}$ instead of $\mathcal{D}(\lambda)$ whenever λ is fixed.

Proof of Lemma 3.9. Take $v \in L^1(S)$, $f \in L^1(V)$, $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ and assume that $\lambda_0 \in \Omega_r$ where the set Ω_r was introduced in the proof of Lemma 3.3.

Notice that by Definition 3.7, of the sets S_n , we can replace S_{n-1} by V in (3.8).

Thus equation (3.8) implies that on V one has

$$u = (\bar{v} - \mathcal{D}f) + \mathcal{D}\mathcal{Q}u. \quad (3.10)$$

We claim that $I - \mathcal{D}\mathcal{Q}$ is an invertible operator.

A simple calculation shows that for $z \in S^c$

$$((\mathcal{D}\mathcal{Q})^n h)(z) = \int_V \int_{S^c} \cdots \int_{S^c} \frac{K(z, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, dz_n) h(z_n)}{(\lambda_0 - d(z)) \prod_{p=1}^{n-1} (\lambda_0 - d(z_p))} \quad (3.11)$$

with $((\mathcal{D}\mathcal{Q})^n h)(z) = 0$ whenever $z \in S$.

As before let $D_n = \{z \in S^c : n_S(z) \geq n\}$, where $n_S(z)$ is the function introduced in Definition 3.5(2), and set $D_0 = S$.

Notice that for $z \in D_m$ when integrating a function over $(z, z_1, \dots, z_n) \in V \times (S^c)^n$ against the kernel $K(z, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, dz_n)$ its integral vanishes outside the domain $D_m \times D_{m-1} \times \cdots \times D_{m-n}$.

Hence, for $n_S(z) < n$ we have $((\mathcal{D}\mathcal{Q})^n h)(z) = 0$. Similarly, if $n_S(z) = n + j$ then

$$((\mathcal{D}\mathcal{Q})^n h)(z) = \int_{D_j} \int_{D_{j+1}} \cdots \int_{D_{n+j-1}} \frac{K(z, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, dz_n) h(z_n)}{(\lambda_0 - d(z)) \prod_{p=1}^{n-1} (\lambda_0 - d(z_p))}.$$

Arguing as in the proof of Lemma 3.5 we obtain

$$\|(\mathcal{D}\mathcal{Q})^n h\|_{L^1(V)} \leq \frac{M^n}{r^n} \mu(D_1) \cdots \mu(D_{n-1}) \mu(S^c) \|h\|_{L^1(V)}.$$

From the convergence of the series $\sum_{n=1}^{\infty} \frac{M^n}{r^n} \mu(D_1) \cdots \mu(D_{n-1})$ (see the proof of Lemma 3.5) the claim follows.

Therefore, from (3.11) we get $u = (I - \mathcal{D}\mathcal{Q})^{-1}(\bar{v} - \mathcal{D}f)$, which implies that

$$\|u\|_{L^1} \leq \max\{1, \|\mathcal{D}\|\} \left(\sum_{n=0}^{\infty} \|(\mathcal{D}\mathcal{Q})^n\| \right) (\|\bar{v}\|_{L^1} + \|f\|_{L^1}).$$

□

Lemma 3.10. *Let V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S be a structural set of type B. Then given $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$, $f \in L^1(V)$ and $v \in L^1(S)$ the following two statements are equivalent:*

- (1) $u = (\bar{v} - \mathcal{D}(\lambda_0) f) + \mathcal{D}(\lambda_0) \mathcal{Q} u$,
- (2) $u = v$ on S and $(\mathcal{A} - \lambda_0 I) u = f$ on S^c .

Proof. For $z \in S$ we have $u(z) = v(z)$ whenever u satisfies either (1) or (2).

For $z \in S^c$, equation

$$u(z) = (\bar{v} - \mathcal{D}(\lambda_0) f)(z) + (\mathcal{D}(\lambda_0) \mathcal{Q} u)(z)$$

is equivalent to

$$u(z) = -\frac{f(z)}{\lambda_0 - d(z)} + \int_V \frac{K(z, dy)}{\lambda_0 - d(z)} u(y)$$

which in turn is equivalent to

$$((\mathcal{A} - \lambda_0 I) u)(z) = f(z).$$

□

Lemma 3.11. *Let V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S be a structural set of type B. Then given $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$, $f \in L^1(V)$ and $u \in L^1(V)$ such that $((\mathcal{A} - \lambda_0 I)u)(z) = f(z)$ for all $z \in S^c$ the following two statements are equivalent:*

- (1) $(\mathcal{R}(\lambda_0) - \lambda_0 I)u_S = f_S$,
- (2) $(\mathcal{A} - \lambda_0 I)u = f$ on V .

Proof. We claim that if for all $z \in S^c$, $((\mathcal{A} - \lambda_0 I)u)(z) = 0$, and $x \in S$ then for all $n \geq n_S(x)$ we have

$$\int_{S^c} u_{S^c}(z) K(x, dz) = \sum_{p=2}^n \int_S u_S(y) K_{S, \lambda_0}^{(p)}(x, dy). \quad (3.12)$$

Let us prove this claim. Since $\mathcal{A}u = \lambda_0 u$ on S^c , we have for all $z \in S^c$

$$d(z) u_{S^c}(z) + \int_V u(w) K(z, dw) = \lambda_0 u_{S^c}(z)$$

which is equivalent to

$$d(z) u_{S^c}(z) + \int_S u_S(y) K(z, dy) + \int_{S^c} u_{S^c}(z') K(z, dz') = \lambda_0 u_{S^c}(z).$$

This in turn is equivalent to

$$u_{S^c}(z) = \frac{1}{\lambda_0 - d(z)} \int_S u_S(y) K(z, dy) + \frac{1}{\lambda_0 - d(z)} \int_{S^c} u_{S^c}(z') K(z, dz'). \quad (3.13)$$

Substituting $u_{S^c}(z')$ by (3.13) in this relation we get

$$\begin{aligned} u_{S^c}(z) &= \frac{1}{\lambda_0 - d(z)} \int_S u_S(y) K(z, dy) + \int_S \int_{S^c} u_S(y) \frac{K(z, dz') K(z', dy)}{(\lambda_0 - d(z)) (\lambda_0 - d(z'))} \\ &\quad + \int_{S^c} \int_{S^c} u_{S^c}(z'') \frac{K(z, dz') K(z', dz'')}{(\lambda_0 - d(z)) (\lambda_0 - d(z'))}. \end{aligned}$$

Given $x \in S$, integrating in $z \in S^c$ w.r.t. $K(x, dz)$ we obtain

$$\begin{aligned} \int_{S^c} u_{S^c}(z) K(x, dz) &= \int_S u_S(y) K_{S, \lambda_0}^{(2)}(x, dy) + \int_S u_S(y) K_{S, \lambda_0}^{(3)}(x, dy) \\ &\quad + \int_{S^c} \int_{S^c} \int_{S^c} u_{S^c}(z'') \frac{K(x, dz) K(z, dz') K(z', dz'')}{(\lambda_0 - d(z)) (\lambda_0 - d(z'))}. \end{aligned}$$

Proceeding inductively, we obtain for all $n \geq 1$ and $x \in S$,

$$\begin{aligned} \int_{S^c} u_{S^c}(z) K(x, dz) &= \sum_{p=2}^n \int_S u_S(y) K_{S, \lambda_0}^{(p)}(x, dy) \\ &\quad + \int_{S^c} \cdots \int_{S^c} u_{S^c}(z_n) \frac{K(x, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, dz_n)}{\prod_{p=1}^{n-1} (\lambda_0 - d(z_p))}. \end{aligned} \quad (3.14)$$

For $n \geq n_S(x)$ the remainder in (3.14) vanishes which proves the claim.

Let us prove (1) $\Rightarrow$ (2). Assuming $\mathcal{R}_S(\lambda_0) u_S = \lambda_0 u_S + f_S$ we have

$$\begin{aligned} f(x) + (\lambda_0 - d(x)) u_S(x) &= \sum_{p=1}^{\infty} \int_S u_S(y) K_{S, \lambda_0}^{(p)}(x, dy) \\ &= \int_S u_S(y) K(x, dy) + \sum_{p=2}^{\infty} \int_S u_S(y) K_{S, \lambda_0}^{(p)}(x, dy) \\ &= \int_S u_S(y) K(x, dy) + \int_{S^c} u_{S^c}(y) K(x, dy) = (\mathcal{Q} u)(x), \end{aligned}$$

where in the last step we use claim (3.12). This proves that $(\mathcal{A} - \lambda_0 I) u = f$ on S . Since we are also assuming that $(\mathcal{A} - \lambda_0 I) u = f$ on S^c , item (2) follows.

Let us now prove (2) $\Rightarrow$ (1). Assuming $(\mathcal{A} u - \lambda_0 I) u = f$ and using the claim (3.12), we have for all $x \in S$,

$$\begin{aligned} f(x) + (\lambda_0 - d(x)) u(x) &= \int_S u_S(y) K(x, dy) + \int_{S^c} u_{S^c}(y) K(x, dy) \\ &= \int_S u_S(y) K(x, dy) + \sum_{p=2}^{\infty} \int_S u_S(y) K_{S, \lambda_0}^{(p)}(x, dy) \\ &= \int_S u_S(y) R_{S, \lambda_0}(x, dy). \end{aligned}$$

This proves that $(\mathcal{R}(\lambda_0) - \lambda_0 I) u_S = f_S$. □

We are now ready for the proof of Theorem 3.8.

Proof of theorem 3.8. Item (3) and the direct implication in (2) are consequences of Lemma 3.11 with $f = 0$.

Next we prove (4) and the converse implication in (2).

Let $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ be an eigenvalue of $\mathcal{R}_S(\lambda_0)$ and $v \in L^1(S)$ be an associated eigenfunction, $\mathcal{R}_S(\lambda_0) v = \lambda_0 v$. Applying Lemma 3.9 with $f = 0$ there exists a function $u \in L^1(V)$ defined recursively by (3.8). Observe that (3.7) is the same as (3.8) when $f = 0$. As

noticed in the proof of Lemma 3.9, (3.8) is equivalent to (3.10), which in this case takes the form $u = \bar{v} + \mathcal{D}Qu$. In turn, by Lemma 3.10, this equation implies that $(\mathcal{A} - \lambda_0 I)u = 0$ on S^c . Now, since $\mathcal{R}_S(\lambda_0)v = \lambda_0 v$ and $(\mathcal{A} - \lambda_0 I)u = 0$ on S^c , by Lemma 3.12, with $f = 0$, we obtain that $(\mathcal{A} - \lambda_0 I)u = 0$ on V .

Finally we prove (1).

Given $\lambda_0 \notin \Sigma_d$, assume that $\lambda_0 \notin \text{sp}(\mathcal{R}_S)$. Then, given $f \in L^1(V)$, there exists $v \in L^1(S)$ such that $(\mathcal{R}_S(\lambda_0) - \lambda_0 I)v = f_S$, where $f_S \in L^1(S)$ stands for the restriction of f to S .

Consider by Lemma 3.9 the function $u \in L^1(V)$ defined recursively by (3.8). As noticed above, (3.8) is equivalent to (3.10), which once more, by Lemma 3.10, is equivalent to $(\mathcal{A} - \lambda_0 I)u = f$ on S^c . Now, since $(\mathcal{R}_S(\lambda_0) - \lambda_0 I)v = f_S$ and $(\mathcal{A} - \lambda_0 I)u = f$ on S^c , by Lemma 3.12 we obtain that $(\mathcal{A} - \lambda_0 I)u = f$ on V .

On the other hand, by (2) this operator must be injective. Therefore $\lambda_0 \notin \text{sp}(\mathcal{A}) \setminus \Sigma_d$. $\square$

Given a structural set S of type B consider the map

$$\Psi_S = \Psi_{S,d,K} : \mathbb{C} \setminus \Sigma_d \rightarrow \mathcal{L}(L^1(V) \times L^1(S), L^1(V))$$

that to each $\lambda \in \mathbb{C} \setminus \Sigma_d$, $f \in L^1(V)$ and $v \in L^1(S)$ associates the unique function $u = \Psi_S(\lambda)(f, v)$ defined recursively by (3.8). Lemma 3.9 proves that this function is well defined.

Remark 3.5. *Given $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$, $f \in L^1(V)$ and $v \in L^1(S)$ such that $(\mathcal{R}(\lambda_0) - \lambda_0 I)v = f_S$ then the function $u = \Psi_S(\lambda_0)(f, v)$ satisfies $(\mathcal{A} - \lambda_0 I)u = f$.*

Proof. By definition $u = \Psi_S(\lambda_0)(f, v)$ satisfies (3.8), which as noticed in the proof of Lemma 3.9 is equivalent to (3.10). Hence, by Lemma 3.10 one has $(\mathcal{A} - \lambda_0 I)u = f$ on S^c . Therefore, by Lemma 3.11 we have that $(\mathcal{A} - \lambda_0 I)u = f$ on V . $\square$

Remark 3.6. *Under the assumptions of Theorem 3.8, the function $\Psi_S : \mathbb{C} \setminus \Sigma_d \rightarrow \mathcal{L}(L^1(V) \times L^1(S), L^1(V))$ is analytic.*

We introduce now the family of *reconstruction operators* $\Phi_S = \Phi_{S,d,K} : \mathbb{C} \setminus \Sigma_d \rightarrow \mathcal{L}(L^1(S), L^1(V))$ defined by

$$\Phi_S(\lambda)(v) := \Psi_S(\lambda)(0, v).$$

Remark 3.7. *If $\lambda \in \text{sp}(\mathcal{R}_S)$ is an eigenvalue of $\mathcal{R}_S(\lambda)$ with eigenfunction $u \in L^1(S)$ then $v = \Phi_S(\lambda)(u)$ is an eigenfunction of $\mathcal{A}$ associated to the same eigenvalue, i.e., $\mathcal{A}v = \lambda v$. This explains the ‘reconstruction’ terminology.*

3.3. Type A isospectral reduction. In this subsection we prove a isospectral reduction theorem for structural sets of type A.

Theorem 3.12. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type A. Let $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ be an eigenvalue of $\mathcal{A}$ and $u \in L^1(V)$ be an associated eigenfunction, $\mathcal{A}u = \lambda_0 u$. Then λ_0 is also an eigenvalue of $\mathcal{R}_S(\lambda_0)$ and $\mathcal{R}_S(\lambda_0) u_S = \lambda_0 u_S$, i.e., u_S is the corresponding eigenfunction for $\mathcal{R}_S(\lambda_0)$.*

Proof. The proof follows the steps of the statement (2) $\Rightarrow$ (1) in Lemma 3.11 with $f = 0$. For each $r > 0$, consider the open set $\Omega_r := \{\lambda \in \mathbb{C} : \text{dist}(\lambda, \Sigma_d) > r\}$, so that $\mathbb{C} \setminus \Sigma_d = \bigcup_{r>0} \Omega_r$. Let $\lambda_0 \in \Omega_r$. We just need to observe that, because S is a structural set of type A, the remainder integral in equality (3.14) converges to 0 as $n \rightarrow +\infty$. Indeed, the equality (3.14) holds regardless of the structural set type. Moreover, for all $x \in S$, the integral

$$\left| \int_{S^c} \cdots \int_{S^c} u_{S^c}(z_n) \frac{K(x, dz_1) K(z_1, dz_2) \cdots K(z_{n-1}, dz_n)}{\prod_{p=1}^{n-1} (\lambda_0 - d(z_p))} \right|$$

is bounded by $\frac{t_n}{r^{n-1}} \|u_{S^c}\|_{L^1}$, and hence converges to 0 as $n \rightarrow +\infty$. □

3.4. Type A quasi-B isospectral reduction. In this subsection we prove a isospectral reduction theorem for structural sets of type A quasi-B.

Let $(E, \|\cdot\|)$ be a Banach space and let $P \in \mathcal{L}(E)$. Let B_1 be the closed unit ball in E . The operator P is called *weakly compact* if the weak closure of PB_1 is compact in the weak topology (see [5]). The set of weakly compact operators is closed in the uniform operator topology of $\mathcal{L}(E)$ ([5, §VI. 4., Corollary 4]).

Lemma 3.13. *Let $\mathcal{A} : E \rightarrow E$ be a weakly compact linear operator on a Banach space E , and $\mathcal{A}_n : E \rightarrow E$ a sequence of linear operators converging to $\mathcal{A}$. Let $u_n \in E$ be a unit eigenvector of $\mathcal{A}_n$ with $\mathcal{A}_n u_n = \lambda_n u_n$ and assume $\lambda_0 = \lim_{n \rightarrow \infty} \lambda_n$ is non zero. Then*

- (1) λ_0 is an eigenvalue of $\mathcal{A}$.
- (2) The sequence $\{u_n\}_n$ is relatively compact.
- (3) Any sublimit u of $\{u_n\}_n$ is an eigenvector of $\mathcal{A}$ with $\mathcal{A}u = \lambda_0 u$.

Proof. By spectrum continuity, $\lambda_0 \in \text{sp}(\mathcal{A})$. Since $\lambda_0 \neq 0$ and $\mathcal{A}$ is weakly compact, λ_0 is an isolated eigenvalue with finite multiplicity. Hence $\text{sp}(\mathcal{A}) = \{\lambda_0\} \cup \Sigma$ for some compact set $\Sigma \subset \mathbb{C}$ with $\lambda_0 \notin \Sigma$. Let $E = F \oplus H$ be the corresponding $\mathcal{A}$ -invariant decomposition where F is the generalized eigenspace associated to λ_0 . Consider a simple closed, positively

oriented curve Γ which isolates λ_0 from Σ and denote by $R(\mathcal{A}, z) := (zI - \mathcal{A})^{-1}$ the resolvent of $\mathcal{A}$. Then the projection $P : E \rightarrow E$ onto F parallel to H is given by

$$P = \frac{1}{2\pi i} \int_{\Gamma} R(\mathcal{A}, z) dz.$$

Of course for large n the operators $\mathcal{A}_n$ admit a similar decomposition of the spectrum

$$\text{sp}(\mathcal{A}_n) = \Lambda_n \cup \Sigma_n,$$

where $\lambda_n \in \Lambda_n$, and Λ_n, Σ_n are closed sets separated by Γ . Hence the operator

$$P_n = \frac{1}{2\pi i} \int_{\Gamma} R(\mathcal{A}_n, z) dz$$

is the projection onto an $\mathcal{A}_n$ -invariant finite dimensional subspace F_n (with same dimension as F). By definition it is clear that $P_n \circ \mathcal{A}_n = \mathcal{A}_n \circ P_n$, which implies that $H_n := \text{Ker}(P_n)$ is also $\mathcal{A}_n$ -invariant. It also follows that P_n converges to P .

Now, since $\lambda_n \in \Lambda_n$, we have $u_n = P_n u_n \in F_n$. The sequence $\tilde{u}_n := P u_n \in F$ is relatively compact because F is finite dimensional. On the other hand, since $P_n \rightarrow P$, we have that $\|u_n - \tilde{u}_n\| = \|P_n u_n - P u_n\| \leq \|P_n - P\|$ converges to 0. Therefore $\{u_n\}_n$ is also relatively compact, which proves (2).

Item (3) is clear. □

It is well known (see e.g., ([6, p. 104]) that integral operators with an uniformly bounded kernel are weakly compact. Therefore, it follows from Proposition 3.1 and Lemma 3.3 that:

Proposition 3.14. *Assume V, K with $d \equiv 0$ satisfy (A1). Then*

- (1) *The operator $\mathcal{A}$ is weakly compact.*
- (2) *Given $\lambda \in \mathbb{C} \setminus \{0\}$, the reduced operators defined by (3.3) are weakly compact.*

Proposition 3.15. *Assume V, K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type A quasi-B. Consider the sequence of kernels K_n in Definition 3.6. Then for every compact set $\Lambda \subset \mathbb{C} \setminus \Sigma_d$,*

$$\lim_{n \rightarrow +\infty} \mathcal{R}_{S,d,K_n}(\lambda) = \mathcal{R}_{S,d,K}(\lambda) \text{ in } \mathcal{L}(L^1(S)),$$

uniformly on $\lambda \in \Lambda$.

Proof. Given $\Lambda \subset \mathbb{C} \setminus \Sigma_d$ compact, choose $r > 0$ so that $\Omega_r = \{\lambda \in \mathbb{C} : \text{dist}(\lambda, \Sigma_d) > r\}$ contains Λ .

Let $K(x, dy) = h(x, y) \mu(dy)$ and $K_n(x, dy) = h_n(x, y) \mu(dy)$. By Definition 3.6 (of type A quasi-B structural sets) $\|h - h_n\|_{\infty} \rightarrow 0$.

We need to compare $R_{S,K,\lambda}^{(p)}$ with $R_{S,K_n,\lambda}^{(p)}$. The density of the first kernel is

$$h_{S,K,\lambda}^{(p)}(x, y) := \int_{S^c} \cdots \int_{S^c} \frac{h(x, z_1) h(z_1, z_2) \cdots h(z_{p-1}, y)}{\prod_{j=1}^{p-1} (\lambda - d(z_j))} \mu(dz_1) \cdots \mu(dz_{p-1}),$$

and a similar formula holds for the density of $R_{S,K_n,\lambda}^{(p)}$ with h_n instead of h .

Write $\hat{\tau}_{K,i} := \sup_{x \in V} \tau_{S,K,i}(x, V)$. Let M be a common upper bound for the density functions $h(x, y)$ and $h_n(x, y)$, with $n \in \mathbb{N}$. Then

$$\|h_{S,K,\lambda}^{(p)} - h_{S,K_n,\lambda}^{(p)}\|_\infty \leq \frac{M^{p-1}}{r^{p-1}} \|h - h_n\|_\infty \sum_{j=1}^{p-1} \hat{\tau}_{K,j} \hat{\tau}_{K_n,p-j}.$$

Then what we need to complete the proof is the following lemma. □

Lemma 3.16. *If τ_n decays super exponentially then so does $\sum_{j=1}^n \tau_j \tau_{n-j}$.*

Proof. It follows from the definition that τ_n decays super exponentially to 0 if and only if for all $L > 0$ there exists $C > 0$ such that $\tau_n \leq C e^{-Ln}$.

Hence, given $L > 0$ there exists $C > 0$ such that $\tau_n \leq C e^{-2Ln}$ for all $n \geq 1$. Therefore

$$\sum_{j=1}^n \tau_j \tau_{n-j} \leq C^2 \sum_{j=1}^n e^{-2Lj} e^{-2L(n-j)} \leq n C^2 e^{2Ln} \leq C^2 e^{-Ln}$$

which proves that the sum $\sum_{j=1}^n \tau_j \tau_{n-j}$ decays super exponentially to 0. □

Proposition 3.17. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type A quasi-B. Consider the sequence of kernels K_n in Definition 3.6. Then given any compact set $\Lambda \subset \mathbb{C} \setminus \Sigma_d$, the following limit exists*

$$\Psi_S(\lambda) = \Psi_{S,d,K}(\lambda) := \lim_{n \rightarrow +\infty} \Psi_{S,d,K_n}(\lambda) \quad (3.15)$$

with uniform convergence in $\lambda \in \Lambda$.

Proof. We claim that for some $m \geq 1$, $\|(\mathcal{D} \mathcal{Q}_K)^m\| < 1$.

From (3.11) in the proof of Lemma 3.9, we have

$$((\mathcal{D} \mathcal{Q}_K)^n h)(z) = \int_{S^c} \cdots \int_{S^c} \frac{K(z, dz_1) \cdots K(z_{n-1}, dz_n) h(z_n)}{(\lambda_0 - d(z)) \prod_{p=1}^{n-1} (\lambda_0 - d(z_p))}$$

for all $z \in S^c$, and $((\mathcal{D} \mathcal{Q}_K)^n h)(z) = 0$ whenever $z \in S$.

Arguing as in the proof of Lemma 3.3 we obtain for all large n

$$\|(\mathcal{D} \mathcal{Q})^n\| \leq \frac{t_n}{r^n} \mu(S^c) \ll 1.$$

From the claim, and since by Definition 3.6(2), $\lim_{n \rightarrow +\infty} \|K - K_n\|_\infty = 0$, we also have $\|(\mathcal{D} \mathcal{Q}_{K_n})^m\| < 1$ for all large enough n . Hence the operators $I - \mathcal{D} \mathcal{Q}_K$ and $I - \mathcal{D} \mathcal{Q}_{K_n}$ are all invertible with uniformly bounded inverses. In particular $\lim_{n \rightarrow +\infty} (I - \mathcal{D} \mathcal{Q}_{K_n})^{-1} = (I - \mathcal{D} \mathcal{Q}_K)^{-1}$.

Given $f \in L^1(V)$ and $v \in L^1(S)$, by the proof of Lemma 3.9, the ‘reconstructed’ function $u_n = \Psi_{S,d,K_n}(\lambda)(f, v)$ is given by

$$u_n = (I - \mathcal{D} \mathcal{Q}_{K_n})^{-1}(\bar{v} - \mathcal{D}f).$$

Therefore $(u_n)_n$ converges in L^1 to $u = (I - \mathcal{D} \mathcal{Q}_K)^{-1}(\bar{v} - \mathcal{D}f)$. $\square$

The previous proposition allows us to define the *limit reconstruction operators* as follows: Given $\lambda \in \mathbb{C} \setminus \Sigma_d$, $\Phi_S(\lambda) : L^1(S) \rightarrow L^1(V)$,

$$\Phi_S(\lambda)(v) := \Psi_S(\lambda)(0, v).$$

Theorem 3.18. *Assume V , K and $d : V \rightarrow \mathbb{C}$ satisfy (A1)-(A2) and S is a structural set of type A quasi-B. Then*

- (1) $\text{sp}(\mathcal{A}) \setminus \Sigma_d \subseteq \text{sp}(\mathcal{R}_S)$.
- (2) *Given $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$, λ_0 is an eigenvalue of $\mathcal{A}$ iff λ_0 is an eigenvalue of $\mathcal{R}_S(\lambda_0)$.*
- (3) *If $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ is an eigenvalue of $\mathcal{A}$ and $u \in L^1(V)$ is an associated eigenfunction, $\mathcal{A}u = \lambda_0 u$, then $\mathcal{R}_S(\lambda_0)u_S = \lambda_0 u_S$, i.e., u_S is the corresponding eigenfunction for $\mathcal{R}_S(\lambda_0)$.*
- (4) *If $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ is an eigenvalue of $\mathcal{R}_S(\lambda_0)$ and v is an associated eigenfunction, $\mathcal{R}_S(\lambda_0)v = \lambda_0 v$, then $u = \Phi_S(\lambda_0)(v)$ is an eigenfunction of $\mathcal{A}$, i.e., $\mathcal{A}u = \lambda_0 u$.*

Proof. Since S is a structural set of type A, by Theorem 3.12, item (3) and the direct implication in (2) follow. The converse implication in (2) will follow from item (4).

Let us prove item (1).

Take $\lambda_0 \notin \text{sp}(\mathcal{R}_S)$. This means that $\mathcal{R}_S(\lambda_0) - \lambda_0 I$ is an invertible operator. We are going to prove that $\mathcal{A} - \lambda_0 I$ is also invertible. By the direct implication in (2) we know that $\mathcal{A} - \lambda_0 I$ is injective. Therefore, it is enough to show that $\mathcal{A} - \lambda_0 I$ is surjective.

To simplify notations we will write $\mathcal{A}$, $\mathcal{A}_n$, $\mathcal{R}$, $\mathcal{R}_n$, Ψ and Ψ_n respectively instead of $\mathcal{A}_{S,d,K}$, $\mathcal{A}_{S,d,K_n}$, $\mathcal{R}_{S,d,K}$, $\mathcal{R}_{S,d,K_n}$, $\Psi_{S,d,K}$ and Ψ_{S,d,K_n} .

Since $\lambda_0 \notin \text{sp}(\mathcal{R})$ and by Proposition 3.15 $\mathcal{R}_n$ converges to $\mathcal{R}$, one has $\lambda_0 \notin \text{sp}(\mathcal{R}_n)$ for all large enough n .

Given $f \in L^1(V)$, because $\mathcal{R}_n(\lambda_0) - \lambda_0 I$ is invertible, there exists $v_n \in L^1(S)$ such that $(\mathcal{R}_n(\lambda_0) - \lambda_0 I)v_n = f_S$. The sequence v_n is bounded because the operator $\mathcal{R}_n(\lambda_0) - \lambda_0 I$ is invertible.

By Proposition 3.14 the operator $\mathcal{R}(\lambda_0)$ can be decomposed as $\mathcal{R}(\lambda_0) = d_S + \widehat{\mathcal{R}}(\lambda_0)$ where d_S is a diagonal operator and $\widehat{\mathcal{R}}(\lambda_0)$ is weakly compact. Analogously, the operator

$\mathcal{R}_n(\lambda_0)$ is decomposed as $\mathcal{R}_n(\lambda_0) = d_S + \widehat{\mathcal{R}}_n(\lambda_0)$, with the same diagonal part d_S and where $\widehat{\mathcal{R}}_n(\lambda_0)$ is also weakly compact. Moreover, $\widehat{\mathcal{R}}_n(\lambda_0)$ converges to $\widehat{\mathcal{R}}(\lambda_0)$ as $n \rightarrow +\infty$.

By weak compactness of $\widehat{\mathcal{R}}(\lambda_0)$, we can assume that $\widehat{\mathcal{R}}(\lambda_0) v_n$ converges to some $w \in L^1(S)$. Since

$$f_S - \widehat{\mathcal{R}}_n(\lambda_0) v_n = (\mathcal{R}_n(\lambda_0) - \lambda_0) v_n - \widehat{\mathcal{R}}_n(\lambda_0) v_n = (d_S - \lambda_0) v_n$$

and

$$\begin{aligned} \|\widehat{\mathcal{R}}_n(\lambda_0) v_n - w\| &\leq \|\widehat{\mathcal{R}}_n(\lambda_0) v_n - \widehat{\mathcal{R}}(\lambda_0) v_n\| + \|\widehat{\mathcal{R}}(\lambda_0) v_n - w\| \\ &\leq \|\widehat{\mathcal{R}}_n(\lambda_0) - \widehat{\mathcal{R}}(\lambda_0)\| \sup_n \|v_n\| + \|\widehat{\mathcal{R}}(\lambda_0) v_n - w\|, \end{aligned}$$

we conclude that $(d_S - \lambda_0) v_n$ converges to $f_S - w$, and hence

$$\lim_{n \rightarrow +\infty} v_n = (d_S - \lambda_0)^{-1}(f_S - w) =: v \text{ in } L^1.$$

By Remark 3.5 the function $u_n = \Psi_n(f, v_n)$ satisfies

$$(\mathcal{A}_n - \lambda_0) u_n = f. \tag{3.16}$$

On the other hand we have

$$\begin{aligned} \|\Psi_n(f, v_n) - \Psi(f, v)\| &\leq \|\Psi_n(f, v_n) - \Psi(f, v_n)\| + \|\Psi(f, v_n) - \Psi(f, v)\| \\ &\leq \|\Psi_n - \Psi\| \sup_n \|v_n\| + \|\Psi\| \|v_n - v\| \end{aligned}$$

which proves that $u_n = \Psi_n(f, v_n)$ converges to $u = \Psi(f, v)$ in L^1 .

Thus, taking the limit in (3.16) we get that $(\mathcal{A} - \lambda_0) u = f$, which proves that $\lambda_0 \notin \text{sp}(\mathcal{A})$.

Finally we prove (4).

Let $\lambda_0 \in \mathbb{C} \setminus \Sigma_d$ be an eigenvalue of $\mathcal{R}(\lambda_0)$ and $v \in L^1(S)$ be an associated eigenfunction, $\mathcal{R}(\lambda_0) v = \lambda_0 v$. Since, by Proposition 3.15, $\mathcal{R}_n$ converges to $\mathcal{R}$, there exist $(\lambda_n)_n$ satisfying $\lambda_0 = \lim_{n \rightarrow +\infty} \lambda_n$ such that $\lambda_n \in \text{sp}(\mathcal{R}_n(\lambda_0))$. By the uniformity of convergence in Proposition 3.15, changing slightly the λ_n if necessary, we may assume that $\lambda_n \in \text{sp}(\mathcal{R}_n(\lambda_n))$.

Let $v_n \in L^1(S)$ be a unit eigenfunction of $\mathcal{R}_n(\lambda_n)$, i.e., $\mathcal{R}_n(\lambda_n) v_n = \lambda_n v_n$. Consider, as before, the weakly compact operators $\widehat{\mathcal{R}}(\lambda_0)$ and $\widehat{\mathcal{R}}_n(\lambda_n)$ so that $\mathcal{R}(\lambda_0)$ and $\mathcal{R}_n(\lambda_n)$ decompose as $\mathcal{R}(\lambda_0) = d_S + \widehat{\mathcal{R}}(\lambda_0)$ and $\mathcal{R}_n(\lambda_n) = d_S + \widehat{\mathcal{R}}_n(\lambda_n)$. Moreover, again by uniformity of convergence, $\widehat{\mathcal{R}}_n(\lambda_n)$ converges to $\widehat{\mathcal{R}}(\lambda_0)$ as $n \rightarrow +\infty$. By Lemma 3.13, extracting a subsequence if necessary we can assume that $(v_n)_n$ converges to v .

Since S is a structural set of type B for K_n , by Theorem 3.8(4), there exists a sequence of eigenfunctions $u_n \in L^1(V)$ such that

$$u_n = \Psi_n(\lambda_n)(0, v_n) \text{ and } \mathcal{A}_n u_n = \lambda_n u_n.$$

Repeating the argument in the proof of item (1), now with $f = 0$, and using uniformity of convergence in Proposition 3.17, we obtain that $u_n = \Psi_n(\lambda_n)(0, v_n)$ converges to $u = \Psi(\lambda_0)(0, v) = \Phi(\lambda_0)(v)$. Hence $\mathcal{A}u = \lambda_0 u$. $\square$

4. ISOSPECTRAL REDUCTION IN INFINITE GRAPHS

In this section we specialize the theory in Section 3 to countably infinite graphs with a finite structural set.

Definition 4.1. *A countable weighted graph is a pair $G = (V, w)$ where V is a countable set and $w : V \times V \rightarrow \mathbb{C}$ is any function, called the weight function of G .*

Definition 4.2. *Let $G = (V, w)$ be a countable weighted graph. Given a positive finite measure μ on V , i.e., $\mu : V \rightarrow [0, +\infty)$ such that $\sum_{v \in V} \mu(v) < +\infty$, we define the Banach space*

$$L^1(V) := \{ f : V \rightarrow \mathbb{C} : \|f\|_1 := \sum_{v \in V} \mu(v) |f(v)| < +\infty \}.$$

Assume $G = (V, w)$ is a countable weighted graph over an infinite set V . The weight function $w : V \times V \rightarrow \mathbb{C}$ determines the following kernel $K(x, \cdot) = \sum_{y \in V} w(x, y) \delta_y(\cdot)$, where δ_y stands for the Dirac measure supported on y .

Let us fix a positive finite reference measure μ on V .

Definition 4.3. *We say that the weight function $w : V \times V \rightarrow \mathbb{C}$ is μ -bounded when there exists $M < +\infty$ such that for all $v, v' \in V$, $|w(v, v')| \leq M \mu(v')$.*

Each μ -bounded function w determines a Markov operator $\mathcal{A}_w : L^1(V) \rightarrow L^1(V)$,

$$(\mathcal{A}_w f)(v) := \sum_{v' \in V} f(v') w(v, v'). \quad (4.1)$$

The following proposition is a simple observation.

Proposition 4.1. *If w is μ -bounded then $\mathcal{A}_w \in \mathcal{L}(L^1(V))$.*

Given a μ -bounded function $w : V \times V \rightarrow \mathbb{C}$, we define

$$\|w\|_\infty := \sup_{v, v' \in V} \frac{|w(v, v')|}{\mu(v')}.$$

Note that identifying the weight function w with the kernel $K(x, \cdot) = \sum_{y \in V} w(x, y) \delta_y(\cdot)$ this norm matches the one defined in 3.5.

Definition 4.4. *Given a countable weighted graph $G = (V, w)$, we associate with it the following point-set map $\varphi_G : V \rightrightarrows V$, $\varphi_G(v) := \{v' \in V : w(v, v') \neq 0\}$.*

Definition 4.5. *Given a countable weighted graph $G = (V, w)$, its support is the set*

$$\Lambda_G := \cap_{j=1}^\infty \varphi_G^j(V).$$

We say that G has finite support when the set $\cap_{j=1}^N \varphi_G^j(V)$ is finite, for some $N \geq 1$. In particular, $|\Lambda_G| < +\infty$.

Definition 4.6. *Given weight functions $w, w' : V \times V \rightarrow \mathbb{C}$, we write $w \ll w'$ if and only if $w'(v, v') = 0$ implies $w(v, v') = 0$, for all $v, v' \in V$.*

Theorem 4.2. *Let $G = (V, w)$ be a countable weighted graph, μ be a positive finite measure μ on V and $S \subset V$ be a finite set. Assume that:*

- (1) w is a μ -bounded weight function on V ;
- (2) S is a structural set of type A for w ;
- (3) $G_n = (V, w_n)$ has finite support for each $n \in \mathbb{N}$;
- (4) $w_n \ll w$, for all $n \in \mathbb{N}$;
- (5) $\lim_{n \rightarrow +\infty} \|w - w_n\|_\infty = 0$.

Then

- (a) $\text{sp}(\mathcal{A}_w) \setminus \{0\} = \text{sp}(\mathcal{R}_{S,w})$.
- (b) *Given $\lambda_0 \in \mathbb{C} \setminus \{0\}$, λ_0 is an eigenvalue of $\mathcal{A}_w$ iff λ_0 is an eigenvalue of $\mathcal{R}_{S,w}(\lambda_0)$.*
- (c) *If $\lambda_0 \in \mathbb{C} \setminus \{0\}$ is an eigenvalue of $\mathcal{A}_w$ and $u \in L^1(V)$ is an associated eigenfunction, $\mathcal{A}_w u = \lambda_0 u$, then $\mathcal{R}_{S,w}(\lambda_0) u_S = \lambda_0 u_S$, i.e., u_S is the corresponding eigenfunction for $\mathcal{R}_{S,w}(\lambda_0)$.*
- (d) *If $\lambda_0 \in \mathbb{C} \setminus \{0\}$ is an eigenvalue of $\mathcal{R}_{S,w}(\lambda_0)$ and v is an associated eigenfunction, $\mathcal{R}_{S,w}(\lambda_0) v = \lambda_0 v$, then $u = \Phi_{S,w}(\lambda_0)(v)$ is an eigenfunction of $\mathcal{A}_w$, i.e., $\mathcal{A}_w u = \lambda_0 u$.*

Proof. This theorem is a corollary of Theorem 3.18. Notice that item (1) implies (A1), while (A2) is automatic since we are taking $d = 0$. We are left to prove that S is a structural set of type A quasi-B for w .

Now, assumption (1) here implies (1) in Definition 3.6, while assumption (5) here implies (2) in Definition 3.6. Hence, we only need to check (3) in Definition 3.6, i.e., that S is a structural set of type B for w_n for all large enough $n \in \mathbb{N}$. We claim that this follows from assumptions (2)-(5) here.

Consider the finite set $V_n := \Lambda_{G_n}$. Since by (3) G_n has finite support we have $V_n = \bigcap_{j=1}^N \varphi_{G_n}^j(V)$ for some large $N \in \mathbb{N}$. Then $\varphi_{G_n}(V_n) \subset V_n$ and $\varphi_{G_n}^N(V) \subset V_n$. For the second inclusion just notice that $\varphi_{G_n}^N(V) \subseteq \varphi_{G_n}^j(V)$ for all $0 \leq j \leq N$.

By assumption (2) there are no cycles in S^c . Hence, because V_n is a finite set, the point-set map $\psi : V_n \rightrightarrows V_n$ defined by $\psi(x) := \varphi_G(x) \cap S^c$ satisfies for some $m \in \mathbb{N}$ and all $x \in V_n$, $\psi^m(x) = \emptyset$.

Let now $\psi_n : V_n \rightrightarrows V_n$ be the point-set map defined by $\psi_n(x) := \varphi_{G_n}(x) \cap S^c$. Items (4) and (5) imply that $\psi_n(x) \subset \psi(x)$ for all $x \in V_n$. Therefore, $\psi_n^m(x) = \emptyset$ for all $x \in V_n$. This together with the inclusion $\varphi_{G_n}^N(V) \subset V_n$ proves that S is a structural set of type B for w_n .

Notice that equality in (a) follows by Remark 3.4. □

We propose now a numerical algorithm to approximate the eigenfunctions of a countably infinite graph. The input and output of the algorithm will consist on the following:

Input:

- a countable weighted graph $G = (V, w)$,
- a finite structural set S ,
- a sequence $(w_n)_n$ of weight functions,
- an integer i ,
- a finite subset V_0 such that $S \subseteq V_0 \subseteq V$,

where $G = (V, w)$, S and $(w_n)_n$ satisfy the assumptions (1)-(5) of Theorem 4.2.

Output: an approximation of the i -th eigenvalue λ of $\mathcal{A}_w$, and an approximation of the values $f(v)$ of a λ -eigenfunction f for $\mathcal{A}_w$ computed at all vertices $v \in V_0$.

Now we describe the steps of the proposed algorithm.

Steps:

- (1) Compute the i -th eigenvalue $\lambda_{i,n}$ of $\mathcal{A}_{w_n}$ for n large, or, alternatively, compute the i -th zero of the analytic function $\det[\mathcal{R}_{S,w_n}(\lambda) - \lambda I] = 0$.
- (2) Compute the associated eigenvector v_0 of the finite dimensional matrix $\mathcal{R}_{S,w_n}(\lambda_{i,n})$, for some large n .

- (3) Use the reconstruction operator $\Psi_{S,w_n}(\lambda_{i,n})(0, v_0)$ to obtain the wanted approximation.

5. AN APPLICATION

In this section we present a concrete application of the method proposed in Section 4, to approximate the eigenfunctions of countably infinite graphs.

Let $G = (\mathbb{N}, w)$ be the countable weighted graph with weight function $w : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{C}$ defined by

- $w(v, v+1) = \frac{1}{2^v}$, for all $v \in \mathbb{N}$;
- $w(v, 1) = 1 - \frac{1}{2^v}$, for all $v \in \mathbb{N}$; and
- $w(v, v') = 0$ otherwise.

Let $w_n : V \times V \rightarrow \mathbb{C}$, $n \in \mathbb{N}$, be a sequence of weight functions defined by

- $w_n(v, v+1) = \frac{1}{2^v}$, for all $v \in \{1, \dots, n\}$;
- $w_n(v, 1) = 1 - \frac{1}{2^v}$, for all $v \in \{1, \dots, n\}$;
- $w_n(m, 1) = 1$, for all $m > n$; and
- $w_n(v, v') = 0$ otherwise.

Consider the finite measure on $\mathbb{N}$ defined by $\mu(v) = \left(\frac{2}{3}\right)^{v-1}$, $v \in \mathbb{N}$.

Let $S = \{1\}$. We have that $G = (\mathbb{N}, w)$, S and the sequence $(w_n)_n$, satisfy the assumptions (1)-(5) of Theorem 4.2. Indeed,

- (1) the weight function w is clearly μ -bounded.
- (2) S is a structural set of type A for w . Since the function $B \mapsto \tau_{S,n,w}(v, B)$ is a measure we just need to observe that for all $n \geq 2$,

- (i) $\tau_{S,n,w}(v, 1) = \left(\prod_{j=0}^{n-2} \frac{1}{2^{v+j}}\right) \left(1 - \frac{1}{2^{v+n-1}}\right) \leq \left(\prod_{j=0}^{n-1} \frac{1}{2^j}\right) \mu(1)$,
- (ii) $\tau_{S,n,w}(v, v+n) = \prod_{j=0}^{n-1} \frac{1}{2^{v+j}} = \left(\prod_{j=0}^{n-1} \frac{1}{2^j}\right) \frac{1}{2^{vn}} \leq \left(\prod_{j=0}^{n-1} \frac{1}{2^j}\right) \mu(v+n)$,
- (iii) $\tau_{S,n,w}(v, v') = 0$ in all other cases.

Therefore, we can take in Definition 3.4 $t_n := \left(\prod_{j=0}^{n-1} \frac{1}{2^j}\right)$, which converges to 0 super exponentially.

- (3) $w_n \ll w$, for all $n \in \mathbb{N}$.
- (4) $G_n = (\mathbb{N}, w_n)$ has finite support for each $n \in \mathbb{N}$ because $\varphi_{G_n}^j(\mathbb{N}) = \{1, \dots, n+1\}$, for all $j \in \mathbb{N}$.

(5) $\lim_{n \rightarrow +\infty} \|w - w_n\|_\infty = 0$. Observe first that

$$\mathcal{A}_w - \mathcal{A}_{w_n} = \begin{bmatrix} 0 & & 0 & & \dots \\ \vdots & \ddots & \vdots & & \\ 0 & & 0 & & \dots \\ -\frac{1}{2^{n+1}} & & 0 & \frac{1}{2^{n+1}} & 0 & \dots \\ -\frac{1}{2^{n+2}} & & 0 & \frac{1}{2^{n+2}} & \frac{1}{2^{n+2}} & \dots \\ -\frac{1}{2^{n+3}} & & & 0 & \frac{1}{2^{n+3}} & \dots \\ & & & & 0 & \dots \\ \vdots & & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Now, since

$$\|w - w_n\|_\infty = \sup_{v, v' \in \mathbb{N}} \frac{|w(v, v') - w_n(v, v')|}{\mu(v')}$$

and $\mu(v) = \left(\frac{2}{3}\right)^{v-1}$, $v \in \mathbb{N}$, a simple calculation shows that $\|w - w_n\|_\infty = \left(\frac{3}{4}\right)^{n+1}$.

We implement now the proposed algorithm with input $i = 2$ and $V_0 = \{1, \dots, 12\}$, to estimate the 2-th eigenvalue λ_2 of $\mathcal{A}_w$, i.e., the eigenvalue, or pair of conjugate eigenvalues, with second largest absolute value, as well as the values of a corresponding eigenfunction computed at all $v \in V_0$.

(1) Let $\lambda_{2,n}$ denote the 2-th eigenvalue of $\mathcal{A}_{w_n}$. Table 1 displays these eigenvalues of $\mathcal{A}_{w_n}$ for $1 \leq n \leq 10$, computed with the help of a computer algebra system.

n	$\lambda_{2,n}$	n	$\lambda_{2,n}$
1	-0.5	6	$-0.139036 \pm 0.162935i$
2	$-0.25 \pm 0.25i$	7	$-0.139075 \pm 0.16291i$
3	$-0.25; -0.125 \pm 0.216506i$	8	$-0.139075 \pm 0.162911i$
4	$-0.125 \pm 0.125i$	9	$-0.139075 \pm 0.162911i$
5	$-0.140218 \pm 0.163328i$	10	$-0.139075 \pm 0.162911i$

TABLE 1. Approximations of the second largest eigenvalue of $\mathcal{A}_w$.

Thus, we can observe that for $n = 8$, the 2-th eigenvalue approximations have stabilized and $\tilde{\lambda}_2 := -0.139075 + 0.162911i$ is an approximation of the 2-th eigenvalue of $\mathcal{A}_w$.

(2) Since for any $n \in \mathbb{N}$ the matrix $\mathcal{R}_{S, w_n}(\tilde{\lambda}_2)$ is a 1×1 -matrix, we can take for the associated eigenvector v_0 any complex number.

- (3) For each $n \in \mathbb{N}$, the reconstruction operator $\Psi_{S,w_n}(\lambda)$ is given by: for $v \in \{2, \dots, n+1\}$,

$$\Psi_{S,w_n}(\lambda)(0, v_0)(v) = \sum_{i=1}^{n-v+2} \lambda^{-i} \left(\prod_{j=0}^{i-2} w_n(v+j, v+j+1) \right) w_n(v+i-1, 1) v_0,$$

and for $v > n+1$,

$$\Psi_{S,w_n}(\lambda)(0, v_0)(v) = \frac{1}{\lambda} v_0.$$

Thus, taking $n = 11$, $\lambda = \tilde{\lambda}_2$ and $v_0 = \tilde{\lambda}_2$, the values of the approximation $(\tilde{f}(v))_{v \in V_0} = \Psi_{S,w_n}(\lambda)(0, v_0)|_{V_0}$ are given in Table 2.

v	$\tilde{f}(v)$	v	$\tilde{f}(v)$
1	$\tilde{\lambda}_2$	7	$0.9685 - 0.0269788i$
2	$0.124679 - 0.253539i$	8	$0.984251 - 0.013679i$
3	$0.513083 - 0.266443i$	9	$0.992126 - 0.00688713i$
4	$0.749921 - 0.175238i$	10	$0.996063 - 0.00345549i$
5	$0.874173 - 0.0990002i$	11	$0.998032 - 0.00173372i$
6	$0.937008 - 0.0524538i$	12	1

TABLE 2. Approximation of an eigenfunction of $\mathcal{A}_w$ associated with λ_2 .

ACKNOWLEDGEMENTS

The first author was supported by “Fundação para a Ciência e a Tecnologia” through the Project UID/MAT/04561/2013.

The second author was partially supported by the Research Centre of Mathematics of the University of Minho with the Portuguese Funds from the “Fundação para a Ciência e a Tecnologia”, through the Project UID/MAT/00013/2013.

REFERENCES

- [1] L. A. Bunimovich and B. Z. Webb, *Isospectral graph reductions and improved estimates of matrices' spectra*, Linear Algebra Appl. **437** (2012), 1429-1457.
- [2] L. A. Bunimovich and B. Z. Webb, *Isospectral graph transformations, spectral equivalence, and global stability of dynamical networks*, Nonlinearity **25** (2012), 211-254.
- [3] L. A. Bunimovich and B. Z. Webb, *Isospectral transformations*, Springer Monographs in Mathematics, Springer, New York, 2014.
- [4] P. Duarte and M. J. Torres, *Eigenvectors of isospectral graph transformations*, Linear Algebra Appl. **474** (2015), 110-123.

- [5] N. Dunford and J. T. Schwartz, *Linear operators, 1 : general theory*, Pure and applied mathematics, Academic press, London, New York, Sydney, 1957.
- [6] L. Weis, *Decompositions of positive operators and some of their applications*, Functional Analysis: Surveys and Recent Results III, North-Holland (1983), 95-115.

CMAF, DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DE LISBOA, CAMPO GRANDE, 1749-016 LISBOA, PORTUGAL.

E-mail address: pedromiguel.duarte@gmail.com

CMAT AND DEPARTAMENTO DE MATEMÁTICA E APLICAÇÕES, UNIVERSIDADE DO MINHO, CAMPUS DE GUALTAR, 4700-057 BRAGA, PORTUGAL

E-mail address: jtorres@math.uminho.pt